



## Guidelines

# PROSPECT guideline for elective caesarean section: updated systematic review and procedure-specific postoperative pain management recommendations

**E. Roofthoof<sup>1,2</sup>, G. P. Joshi,<sup>3</sup> N. Rawal,<sup>4</sup> M. Van de Velde,<sup>5</sup> and on behalf of the PROSPECT Working Group\* of the European Society of Regional Anaesthesia and Pain Therapy and supported by the Obstetric Anaesthetists' Association**

1 Consultant, Department of Anesthesiology, GZA Sint-Augustinus Hospital, Antwerp, Belgium

2 PhD Student, 5 Professor and Chair, Department of Cardiovascular Sciences, KULeuven and UZLeuven, Leuven, Belgium

3 Professor, Department of Anesthesiology and Pain Management, University of Texas Southwestern Medical Center, Dallas, Texas, USA

4 Professor, Department of Anesthesiology, Orebro University, Orebro, Sweden

## Summary

Caesarean section is associated with moderate-to-severe postoperative pain, which can influence postoperative recovery and patient satisfaction as well as breastfeeding success and mother-child bonding. The aim of this systematic review was to update the available literature and develop recommendations for optimal pain management after elective caesarean section under neuraxial anaesthesia. A systematic review utilising procedure-specific postoperative pain management (PROSPECT) methodology was undertaken. Randomised controlled trials published in the English language between 1 May 2014 and 22 October 2020 evaluating the effects of analgesic, anaesthetic and surgical interventions were retrieved from MEDLINE, Embase and Cochrane databases. Studies evaluating pain management for emergency or unplanned operative deliveries or caesarean section performed under general anaesthesia were excluded. A total of 145 studies met the inclusion criteria. For patients undergoing elective caesarean section performed under neuraxial anaesthesia, recommendations include intrathecal morphine 50–100 µg or diamorphine 300 µg administered pre-operatively; paracetamol; non-steroidal anti-inflammatory drugs; and intravenous dexamethasone administered after delivery. If intrathecal opioid was not administered, single-injection local anaesthetic wound infiltration; continuous wound local anaesthetic infusion; and/or fascial plane blocks such as transversus abdominis plane or quadratus lumborum blocks are recommended. The postoperative regimen should include regular paracetamol and non-steroidal anti-inflammatory drugs with opioids used for rescue. The surgical technique should include a Joel-Cohen incision; non-closure of the peritoneum; and abdominal binders. Transcutaneous electrical nerve stimulation could be used as analgesic adjunct. Some of the interventions, although effective, carry risks, and consequentially were omitted from the recommendations. Some interventions were not recommended due to insufficient, inconsistent or lack of evidence. Of note, these recommendations may not be applicable to unplanned deliveries or caesarean section performed under general anaesthesia.

Correspondence to: M. Van de Velde

Email: marc.vandavelde@uzleuven.be

Accepted: 5 November 2020

Re-use of this article is permitted in accordance with the Creative Commons Deed, Attribution 2.5, which does not permit commercial exploitation.

Keywords: analgesia; caesarean section; caesarean delivery; pain

Twitter: @MarcVandeVelde6

\* Members of the PROSPECT Working Group, see Appendix 1.; @MarcVandeVelde6

## Recommendations

- 1 Implement strategies to minimise systemic opioid utilisation and develop individualised or stratified post-discharge opioid prescribing practices to reduce unnecessary opioid analgesic consumption after elective caesarean section.
- 2 Add intrathecal morphine 50–100 µg or diamorphine 300 µg to spinal anaesthesia. Epidural morphine 2–3 mg or diamorphine 2–3 mg may be used as an alternative, for example, when an epidural catheter is used as part of a combined spinal-epidural technique.
- 3 Prescribe paracetamol and a non-steroidal anti-inflammatory drug (NSAID) administered after delivery and continued regularly postoperatively.
- 4 Administer a single dose of intravenous (i.v.) dexamethasone after delivery in the absence of contraindications.
- 5 Consider a single injection of local anaesthetic infiltration, continuous wound local anaesthetic infusion and/or fascial plane blocks, if intrathecal morphine is not used.
- 6 Use a surgical technique that includes the Joel-Cohen incision, non-closure of the peritoneum and abdominal binders.
- 7 Consider the use of transcutaneous electrical nerve stimulation as an analgesic adjunct.

## Why was this guideline developed?

Caesarean section is associated with moderate-to-severe postoperative pain which may influence recovery, psychological maternal well-being, breastfeeding and mother-child bonding. The aim of this guideline is to provide clinicians with updated evidence for optimal pain management following elective caesarean section under neuraxial anaesthesia.

## What other guidelines are available on this topic?

The procedure-specific postoperative pain management (PROSPECT) recommendations for pain management after caesarean section were published in 2014; however, an update assessing analgesic interventions was necessary given developments in clinical practice. The American College of Obstetricians and Gynecologists has provided

recommendations for postpartum pain management which are available on their website.

## How does this guideline differ from other guidelines?

The updated systematic review further confirms the previous recommendations. Also, an updated PROSPECT approach was used to develop the current recommendations such that the available evidence is critically assessed for current clinical relevance and the use of simple, non-opioid analgesics such as paracetamol and NSAIDs as basic analgesics. This approach reports true clinical effectiveness by balancing the invasiveness of the analgesic interventions and the degree of pain after surgery, as well as balancing efficacy and adverse effects.

## Introduction

Caesarean section is associated with moderate-to-severe postoperative pain in a significant proportion of women, which may delay recovery and return to activities of daily living; impair mother-child bonding; impact maternal psychological well-being; and may complicate breastfeeding [1]. Furthermore, inadequate postoperative pain relief may lead to hyperalgesia and persistent postoperative pain [2].

Pain after caesarean section is often under-treated due to unfounded fears that analgesic drugs or interventions might induce maternal and neonatal side-effects and because the severity of post-caesarean section pain is often underestimated [3]. Based on a systematic review performed in 2014 [4], the PROSPECT Working Group [5,6], which is a collaboration of surgeons and anaesthetists, previously provided recommendations for pain management in women undergoing caesarean section. Recently, several new techniques have been developed to manage pain after caesarean, such as the quadratus lumborum block; slow-release local anaesthetics; and non-pharmacological approaches. Additionally, in the last decade, attention has shifted to reduce opioid use and to implement protocols for enhanced recovery after caesarean section. Therefore, an updated systematic review on analgesic interventions for pain management after elective caesarean section performed using neuraxial anaesthesia

was needed. In addition, it was deemed necessary to re-assess the recommendations to align them with the updated PROSPECT approach that considers current clinical relevance and clinical effectiveness by balancing the invasiveness of the analgesic interventions and the degree of pain after surgery, as well as balancing efficacy and adverse effects [7,8].

The aim of this systematic review was to provide updated recommendations based on recent literature assessing the impact of analgesic and surgical approaches on pain after elective caesarean section performed under neuraxial anaesthesia. Postoperative pain scores were the primary outcome measures. Other recovery outcomes assessed included cumulative opioid consumption and adverse effects. The limitations of the available evidence were also assessed. The ultimate aim was to develop recommendations for pain management after elective caesarean section performed under neuraxial anaesthesia.

## Methods

The methods of this review adhered to the previously reported PROSPECT methodology [8]. Specific to this study, the Embase, MEDLINE, PubMed and Cochrane databases (Cochrane Central Register of Controlled Trials; Cochrane Database of Abstracts or Reviews of Effects; Cochrane Database of Systematic Reviews) were searched for randomised controlled trials, systematic reviews and meta-analyses published between 1 May 2014 and 22 October 2020. The search terms used were: (cesarean section OR cesarean OR cesarean delivery) AND (pain OR postoperative pain OR analgesia OR anesthesia OR anaesthesia OR anesthetic) AND (anesthetics neuraxial OR intrathecal OR spinal OR epidural analgesia OR paravertebral blocks OR peripheral nerve OR peripheral block OR regional nerve OR transversus abdominis plane block OR infiltration OR instillation OR NSAID OR COX-2 OR paracetamol OR acetaminophen OR gabapentin OR pregabalin OR clonidine OR opioid OR ketamine OR corticosteroid OR dexamethasone OR peritoneal closure OR skin incision OR skin closure). Only studies in which patients underwent elective caesarean section under neuraxial anaesthesia were included.

Quality assessment, data extraction and data analysis adhered to the PROSPECT methodology [8]. In this study, we defined a change of more than 10/100 mm on the visual analogue scale or numerical rating score as clinically-relevant [8]. The effectiveness of each intervention for each outcome was evaluated qualitatively by assessing the number of studies showing a significant difference between

treatment arms. We also evaluated for each study if patients received 'basic' analgesia (i.e. paracetamol and/or NSAIDs) and 'baseline' analgesia (i.e. routine administration of an analgesic additional to the study intervention). We decided not to perform a meta-analysis a priori due to heterogeneity in study design and result reporting, restricting pooled analysis.

Recommendations were made according to PROSPECT methodology [8]. In brief, this involved a grading of A–D according to the overall level of evidence, as determined by the quality of studies included, consistency of evidence and study design. The proposed recommendations were sent to the PROSPECT Working Group for review and comments and a modified Delphi approach was utilised as previously described. Once a consensus was achieved the lead authors drafted the final document, which was ultimately approved by the Working Group. The Obstetric Anaesthetists' Association Executive Committee were consulted on the final PROSPECT recommendations and offered their support.


## Results

A total of 145 studies were included, of which 126 were randomised controlled trials and 19 were systematic reviews and meta-analyses (Fig. 1) [9–153]. The methodological quality assessments of the 126 randomised controlled trials included in the final qualitative analysis are summarised in online Supporting Information Table S1. The characteristics of the included studies are shown in online Supporting Information Tables S2 and S3.

### **Systemic non-opioid and opioid analgesics**

When paracetamol was administered pre-operatively rather than at the end of surgery, only minor differences were noted [9]. In one study, rectal paracetamol was shown to be superior to pre-operative oral paracetamol combined with i.v. paracetamol at the end of surgery [10]. In one study, opioid consumption was reduced with i.v. paracetamol compared with placebo but there was no difference in pain scores [11]. In another study, no differences in opioid consumption and pain scores were noted with i.v. paracetamol [12].

A meta-analysis concluded that systemic NSAIDs reduced pain scores, decreased opioid consumption, reduced opioid-related side-effects and increased patient satisfaction [13]. A Cochrane review evaluated oral analgesics, comprising primarily but not exclusively NSAIDs, but could not draw any conclusions due to the low quality of studies, small number of included patients and substantial heterogeneity in the studied drugs



**Figure 1** Flow diagram of studies included in this systematic review.

(paracetamol; celecoxib; ibuprofen; gabapentin; combination) [14]. Inthigood et al. evaluated a single dose of i.v. parecoxib 40 mg and noted better pain scores than with placebo [15]. Three studies compared an NSAID with an opioid and demonstrated equally effective or superior analgesia with NSAIDs [16–18]. The addition of rectal diclofenac to pentazocine was also associated with better analgesia than pentazocine alone [19].

Four randomised controlled trials [20–23] and a meta-analysis [24] evaluated pre-operative gabapentinoids for analgesia after caesarean section. No significant benefits were reported with gabapentin when added to a multimodal

analgesia regimen [20]. The multimodal regimen consisted of intrathecal morphine, rectal and oral paracetamol and i.v. and oral NSAID [20]. Administration of pregabalin combined with intramuscular diclofenac, but without intrathecal morphine, was associated with lower pain scores and reduced opioid requirements [21]. In another study, gabapentin provided superior analgesia compared with intrathecal fentanyl [22]. In the two latter studies, basic analgesia consisted of diclofenac [21,22]. In a study conducted on patients who did not receive any basic analgesia, adding vitamin B complex to gabapentin reduced pain scores and opioid consumption compared with the use of gabapentin alone [23]. A systematic review

reported a clinically significant reduction in 24-h pain scores with pre-operative gabapentin. Side-effects such as sedation and dizziness were reported in several of the included studies [24].

Adding i.v. lidocaine to i.v. patient-controlled analgesia (PCA) with morphine did not improve pain scores or opioid consumption [25]. One randomised controlled trial evaluated the effects of i.v. ketamine on postoperative analgesia [26]. A bolus of i.v. ketamine after delivery of the fetus reduced pain and rescue analgesics in the first 12 h after caesarean section [26]. In the latter study, no basic analgesia or additional baseline analgesia was given [26]. A meta-analysis on the i.v. use of ketamine demonstrated marginal improvements in pain scores and a mild reduction in morphine consumption [27].

Compared with sufentanil PCA alone, the addition of dexmedetomidine to a sufentanil PCA in the postoperative period was associated with lower pain scores, reduced sufentanil consumption, reduced need for rescue analgesia and a higher patient satisfaction. However, the improved pain scores were not clinically relevant [28]. In the latter study, no basic or additional baseline analgesia was given [28].

Four randomised controlled trials evaluated the use of i.v. dexamethasone [29–32]. Use of i.v. dexamethasone was associated with better pain scores; prolongation of analgesic effect [29]; a reduction in opioid consumption [30]; and a reduced need for postoperative anti-emetics [31]. One study reported better analgesia when dexamethasone was administered as wound infiltration as opposed to i.v. administration [32]. Intravenous dexamethasone was not as effective as i.v. tramadol [32].

Several studies compared various systemic opioids (oxycodone; sufentanil; tramadol; dezocine; butorphanol; hydromorphone; tapentadol) [33–40]. No individual drug was clearly superior in terms of analgesia or side-effect profile compared with any other opioid.

### **Neuraxial adjuvant drugs**

One meta-analysis [41] and three randomised controlled trials [42–44] evaluated the administration of intrathecal morphine. The meta-analysis compared low (50–100 µg) and high (> 100 µg) doses of intrathecal morphine and concluded that high doses increase the duration of analgesia but were more likely to be associated with side-effects. Pain scores were similar in both groups [41]. A dose-response study of intrathecal morphine showed that 50 µg doses were as effective as 100 µg and 150 µg, with a similar requirement for rescue opioids. The risk of pruritus was lowest after 50 µg morphine [42]. In patients with an anticipated high pain intensity (such as patients with chronic

pelvic pain), pain scores with movement were lower in patients receiving 300 µg vs. those receiving 150 µg intrathecal morphine [43]. In a comparative study, intrathecal morphine provided better analgesia compared with epidural morphine and patient-controlled epidural analgesia of ropivacaine with sufentanil [44]. In two studies, women were offered to choose the analgesic strategy and select either no intrathecal morphine or a low or high dose of intrathecal morphine [45,46]. Having a choice did not impact on rescue opioid consumption, but women were very good in predicting their actual opioid needs. Choosing high-dose intrathecal morphine was associated with increased rescue analgesia and more vomiting [45,46]. Apart from one study [44], all studies used basic analgesia with NSAIDs [42,43,46] or a combination of NSAIDs and paracetamol [45]. Intrathecal morphine was similar to intrathecal hydromorphone in a recent trial by Sharpe et al. [47].

Ten trials evaluated the neuraxial administration of α<sub>2</sub>-agonists such as clonidine and dexmedetomidine [48–57]. A meta-analysis showed that neuraxial clonidine increased the duration and quality of analgesia and reduced morphine consumption [48]. However, more side-effects such as hypotension and intra-operative sedation were noted. No improvements in analgesia were reported with intrathecal or i.v. clonidine, whether administered alone [49] or in combination with intrathecal morphine [50]. One study demonstrated the superiority of intrathecal clonidine to intrathecal fentanyl [51]. Addition of epidural dexmedetomidine to combined spinal-epidural anaesthesia resulted in improved intra-operative and postoperative analgesia and less requirements for opioid rescue [52]. A comparison of intrathecal dexmedetomidine with intrathecal morphine did not demonstrate any significant differences in duration of analgesia, pain scores or need for rescue analgesia. However, both intrathecal morphine and intrathecal dexmedetomidine provided better analgesia when compared with isobaric bupivacaine (53). Administration of intrathecal dexmedetomidine resulted in improved postoperative analgesia when compared with isobaric bupivacaine or ropivacaine alone [54,55]. Intrathecal dexmedetomidine combined with intrathecal magnesium sulphate or intrathecal morphine improved analgesia which was of longer duration than analgesia produced by magnesium sulphate alone [56,57]. Adding intrathecal fentanyl to bupivacaine improved initial analgesia [58]. However, when morphine is also added to the intrathecal mixture, fentanyl might induce acute opioid tolerance and result in greater opioid consumption [59]. Intrathecal buprenorphine [60] and epidural hydromorphone [61] resulted in improved postoperative

analgesia and reduced opioid consumption compared with intrathecal bupivacaine or ropivacaine alone.

A meta-analysis evaluating the effect of neuraxial magnesium on postoperative analgesia demonstrated a longer duration of sensory block, lower pain scores and reduced rescue analgesia requirements than neuraxial mixtures of local anaesthetic without magnesium [62].

The use of intrathecal midazolam was evaluated in several studies [63,64]. A comparative study demonstrated that intrathecal magnesium and intrathecal sufentanil were superior to intrathecal midazolam [63]. Intrathecal midazolam prolonged the duration of spinal anaesthesia when compared with placebo [64]. Intrathecal ketamine prolonged analgesia when compared with fentanyl [65,66]. A meta-analysis showed that intrathecal neostigmine improved analgesia after caesarean section, although it was associated with an increased risk of nausea and vomiting [67]. A study showed that a faster speed of intrathecal injection of fentanyl and local anaesthetic results in improved postoperative analgesia with a more sustained duration [68].

#### **Local and regional analgesia techniques**

Intraperitoneal local anaesthetic instillation resulted in lower early pain scores [69], and reduced pain scores at 24 h in a sub-group in which the peritoneum was closed [69]. The use of topical analgesia (e.g. eutectic mixture of local anaesthetic cream) failed to reduce pain scores at 24 and 48 h [70].

Three studies demonstrated that local anaesthetic wound infiltration reduced pain scores and the need for rescue analgesia during the first 24 h after caesarean section [71–73], while one study showed only limited benefits [74]. Apart from one study [72], basic analgesia with ibuprofen and paracetamol was provided. Another two studies which used multimodal analgesia showed improved pain scores, less morphine consumption and higher breastfeeding comfort with continuous wound infusion compared with no infusion [75,76]. Local anaesthetic wound infusion resulted in similar analgesic effects as intrathecal morphine [76]. A meta-analysis confirmed that both single-shot local anaesthetic wound infiltration and continuous wound infusion reduce postoperative opioid consumption and mildly improve pain scores [77]. Pain scores were similar whether the catheter was placed preperitoneal or subcutaneously [78]. Adding ketorolac improved analgesia of wound infiltration and reduced opioid consumption [79]. In a recent study, ketorolac added to wound infiltration did not improve analgesia but intrathecal morphine was administered in both groups [80]. Magnesium and

dexmedetomidine as adjuvants to wound infiltration reduced pain scores [81,82].

A rectus sheath block provides no additional analgesic benefit when added to multimodal analgesia which also includes intrathecal morphine [83]. Adding a field block after caesarean section to intrathecal morphine also did not improve analgesia after caesarean section [84].

There were five studies that compared transversus abdominis plane (TAP) blocks against placebo or no TAP block [85–89]. Apart from one study [85], all studies noted that TAP blocks improved pain relief, increased patient satisfaction and resulted in a reduction of rescue analgesia. A comparison between lateral and posterior approaches concluded that the posterior approach resulted in better pain scores which was only clinically relevant at 12 h postoperatively. This approach also resulted in reduced need for rescue analgesia [90]. Comparison between surgeon-administered and anaesthetist-administered TAP blocks did not show any differences in postoperative analgesia [91].

Several studies evaluated the role of local anaesthetic adjuvants for TAP blocks. Pain scores, opioid consumption and duration of analgesia were significantly improved when dexamethasone was added to local anaesthetic for TAP blocks [92]. Fentanyl added to TAP blocks failed to improve the quality of analgesia [93]. The addition of  $\alpha_2$ -agonists (clonidine or dexmedetomidine) prolonged the duration of analgesia, reduced the need for rescue drugs and improved patient satisfaction [94–96]. However, mild sedation was noted in some patients [94–96].

Several studies compared TAP blocks with alternative regional anaesthesia techniques [97–103]. In a comparison of TAP blocks with epidural analgesia which included high-dose epidural morphine, improved analgesia with the epidural analgesia was noted [97]. Three studies compared intrathecal morphine with TAP blocks [98–100]. In two of these, there was better analgesia with intrathecal morphine and a reduced requirement for rescue analgesia. However, postoperative mobilisation and return of gastro-intestinal function was better with TAP blocks [98,99]. The third study could not discriminate between the two techniques in terms of pain relief and other clinical outcomes [100]. Three randomised controlled trials compared TAP blocks with continuous local anaesthetic wound infusion and noted no differences in postoperative analgesia [101–103].

Three meta-analyses confirmed the efficacy of TAP blocks for analgesia after caesarean section but concluded that they do not confer any benefit over intrathecal morphine [104–106]. A combination of ilioinguinal and iliohypogastric nerves block with TAP blocks vs. no blocks

resulted in less rescue opioid consumption and lower pain scores [107]. Adding dexmedetomidine to a ropivacaine bilateral ultrasound-guided TAP block resulted in lower postoperative pain scores and less rescue opioid [108]. Quadratus lumborum blocks were evaluated in 11 trials [109–119]. Compared with a sham block, quadratus lumborum blocks produced better analgesia. In two trials quadratus lumborum blocks were found to be superior to a TAP blocks [113,114,118,119]. In one study, quadratus lumborum blocks were less effective than a single epidural bolus of local anaesthetic [115]. Adding quadratus lumborum blocks to intrathecal morphine did not improve analgesia [116]. However, in a direct comparison, a quadratus lumborum block was similar to intrathecal morphine [117]. Two recent meta-analyses evaluated TAP blocks, wound infusion and quadratus lumborum blocks with or without intrathecal morphine and concluded that all three regional anaesthetic techniques are superior to no regional technique in the absence of intrathecal morphine [120,121]. When intrathecal morphine is administered, adding these techniques confers no further advantages.

Two studies recently evaluated the erector spinae plane block (ESP) compared with TAP block and intrathecal morphine and in both studies the ESP block improved analgesia [122,123].

### **Patient-controlled epidural analgesia**

Patient-controlled epidural analgesia added to intrathecal morphine resulted in a further lowering of postoperative pain scores and less need for rescue opioid [124]. Adding fentanyl to patient-controlled epidural analgesia with levobupivacaine did not improve analgesia [125].

### **Postoperative interventions**

Several investigators reported on the beneficial effects of transcutaneous electrical nerve stimulation on pain scores, rescue analgesia use and patient satisfaction [126,127]. A study demonstrated that self-administered oral opioid analgesia was as effective as parenteral nurse-administered drugs [128]. A comparison of a fixed time-interval with on-demand oral analgesia concluded that the latter was associated with better pain scores [129]. One study evaluated the use of relaxation sounds intra- or postoperatively and showed improved pain scores [130]. One study evaluated the use of early skin-to-skin contact between mother and baby and noted no differences in postoperative pain scores [131].

Three studies evaluated the use of elastic abdominal binders after caesarean section [132–134]. In all three, a clinically-relevant reduction in pain scores and rescue

analgesia was noted [132–134]. Two studies evaluated the application of manual cervical dilation at the end of caesarean section and compared it with no cervical dilation and came to conflicting conclusions [135,136]. One study noted improved pain scores until 7 days postoperatively [135] while the other did not report any pain reduction [36]. Pre-operative vaginal cleansing resulted in minor but statistically significant reductions in postoperative pain scores [137].

### **Surgical interventions**

A systematic review [138] confirmed the superiority of the Joel-Cohen (also called modified Misgav-Ladach) incision compared with Pfannenstiel incision in reducing postoperative pain [139]. No differences in pain scores were noted between using a scalpel vs. diathermy for the skin incision [139].

A blunt fascial opening resulted in less postoperative pain [140]. The older technique of extraperitoneal section was associated with better pain scores up to 48 h postoperatively [141]. In one study, the absence of making a bladder flap at opening the uterus resulted in clinically-relevant improvements in postoperative pain scores [142]. A comparison between uterine exteriorisation and in situ closure of the uterus showed more postoperative pain with exteriorised uteri [143]. However, one meta-analysis did not show any difference in postoperative pain between the two modalities of uterine closure [144].

A comparison between two techniques of pyramidalis muscle dissection found no differences in postoperative pain [145]. Reduced pain scores when the peritoneum was not closed were reported [146]. One study reported a significant reduction in postoperative pain scores when the rectus muscle was not re-approximated [147]. A Cochrane review noted minimal evidence for reduced pain scores when the peritoneum was not closed after caesarean section [148]. When applying laser irradiation to the caesarean section wound at the end of surgery, less pain during the first 24 h postoperatively was noted [149,150]. No differences in pain scores were noted between interrupted and continuous wound suturing [151]. Similarly, two meta-analyses did not show any difference in pain scores whether skin closure was performed with sutures or staples [152,153].

## **Discussion**

The majority of the studies included in this systematic review were determined to be of high quality. The updated literature strengthens the previous PROSPECT recommendations for pain management in patients

**Table 1** Overall recommendations for pain management in patients undergoing elective caesarean section.

<b>Pre-operatively</b>
<ul style="list-style-type: none"> <li>Intrathecal long-acting opioid (e.g. morphine 50–100 µg or diamorphine up to 300 µg) (Grade A). Epidural morphine 2–3 mg or diamorphine up to 2–3 mg may be used as an alternative, for example, when an epidural catheter is used as part of a combined spinal-epidural technique (Grade A)</li> <li>Oral paracetamol (Grade A)</li> </ul>
<b>Intra-operative after delivery</b>
<ul style="list-style-type: none"> <li>Intravenous paracetamol if not administered pre-operatively (Grade A)</li> <li>Intravenous non-steroidal anti-inflammatory drugs (Grade A)</li> <li>Intravenous dexamethasone (Grade A)</li> <li>If intrathecal morphine not used, local anaesthetic wound infiltration (single-shot) or continuous wound infusion and/or regional analgesia techniques (fascial plane blocks such as transversus abdominis plane blocks and quadratus lumborum blocks) (Grade A)</li> </ul>
<b>Postoperative</b>
<ul style="list-style-type: none"> <li>Oral or intravenous paracetamol (Grade A)</li> <li>Oral or intravenous non-steroidal anti-inflammatory drugs (Grade A)</li> <li>Opioid for rescue or when other recommended strategies are not possible (e.g. contra-indications to regional anaesthesia) (Grade D)</li> <li>Analgesic adjuncts include transcutaneous electrical nerve stimulation (Grade A)</li> </ul>
<b>Surgical technique</b>
<ul style="list-style-type: none"> <li>Joel-Cohen incision (Grade A)</li> <li>Non-closure of peritoneum (Grade A)</li> <li>Abdominal binders (Grade A)</li> </ul>

undergoing elective caesarean section and modifies it in certain aspects. The updated PROSPECT methodology further strengthens the recommendations, because it goes beyond assessment of the available evidence based solely on statistical analysis [8].

Of note, it is essential to highlight that this guideline focuses on elective caesarean section under neuraxial anaesthesia. Importantly, these recommendations should not be applied to other patient populations such as emergency or unplanned caesarean section or surgery performed under general anaesthesia.

The recommended strategies have sufficient procedure-specific evidence and have a positive balance of clinical benefits and risk of side-effects (Table 1). Basic analgesia after caesarean section should always consist of paracetamol and NSAIDs started intra-operatively (after delivery) and continued postoperatively, unless there are contra-indications. Of note, several studies demonstrated equally good pain control with NSAIDs compared with opioids. Regular administration of basic analgesics is important to limit the need for rescue opioid analgesia. Moreover, studies investigating an analgesic strategy to manage pain relief after caesarean section should not omit this basic strategy of analgesia so as to establish the additional value of an investigational approach. In addition to basic analgesics, i.v. dexamethasone demonstrated positive effects on pain scores and opioid consumption. In

addition, i.v. dexamethasone provides anti-emetic prophylaxis. Thus, i.v. dexamethasone is recommended. Caution is required in patients with glucose intolerance.

Intrathecal morphine at doses of 100 µg or lower is recommended. Doses lower than 100 µg result in adequate analgesia with a reduced incidence of side-effects. Recently, Sharawi et al. confirmed the safety of intrathecal morphine when used in patients undergoing caesarean section [154]. Importantly, basic analgesics (i.e. paracetamol and NSAIDs) and i.v. dexamethasone should be used with intrathecal morphine. Of note, the National Institute of Health and Care Excellence (NICE) guidelines in the UK recommend intrathecal diamorphine as an alternative to intrathecal morphine [155]. Intrathecal diamorphine 300 µg is recommended. When spinal anaesthesia is not possible or when an epidural catheter is in situ, epidural morphine or diamorphine both in doses of 2–3 mg can be used.

Various local anaesthetic techniques such as TAP blocks, quadratus lumborum blocks and local anaesthetic wound infiltration are effective in reducing pain scores and opioid requirements. Given that the potential side-effects of these regional analgesic techniques are limited, they are recommended. However, the additional value of any of these techniques when combined with intrathecal morphine appears to be minimal. Therefore, these blocks may be administered if intrathecal morphine is not used.

**Table 2** Analgesic interventions that are not recommended for pain management in patients undergoing elective caesarean section.

	<b>Intervention</b>	<b>Reason for not recommending</b>
<b>Pre-operative</b>	Gabapentinoids	Limited procedure-specific evidence and concerns of side-effects
<b>Intra-operative</b>	Intravenous ketamine	Limited procedure-specific evidence and concerns of side-effects
	Intravenous dexmedetomidine	Limited procedure-specific evidence and concerns of side-effects
	Intravenous tramadol and butorphanol	Limited procedure-specific evidence
	Neuraxial clonidine	Inconsistent procedure-specific evidence and concerns of side-effects
	Neuraxial dexmedetomidine	Inconsistent procedure-specific evidence and concerns for side-effects
	Intrathecal buprenorphine	Limited procedure-specific evidence
	Epidural hydromorphone	Limited procedure-specific evidence
	Intrathecal midazolam	Limited procedure-specific evidence and concerns of side-effects
	Intrathecal neostigmine	Concerns of side-effects
	Intrathecal ketamine	Limited procedure-specific evidence and concerns of side-effects
	Intraperitoneal local anaesthetic	Lack of procedure-specific evidence
	Topical skin analgesia	Lack of procedure-specific evidence
	Clonidine added to TAP	Lack of procedure-specific evidence
	Dexmedetomidine added to TAP	Limited procedure-specific evidence
	Fentanyl added to TAP	Lack of procedure-specific evidence
	Rectus sheath block	Lack of procedure-specific evidence
Field block	Lack of procedure-specific evidence	
Music	Limited procedure-specific evidence	
<b>Postoperative</b>	Skin-to-skin contact	Limited procedure-specific evidence
	Intravenous lidocaine	Lack of procedure-specific evidence
	Patient controlled epidural analgesia	Limited procedure-specific evidence and concerns of side-effects
<b>Surgical technique</b>	Method of incision: diathermy	Inconsistent procedure-specific evidence
	Absence of a bladder flap	Limited procedure-specific evidence
	Blunt fascial opening	Limited procedure-specific evidence
	Uterine exteriorisation	Inconsistent procedure-specific evidence
	Skin incision lasering postoperatively	Limited procedure-specific evidence
	Type of skin closure	Lack of procedure-specific evidence
	Vaginal cleansing	Lack of procedure-specific evidence
	Cervical dilation	Inconsistent procedure-specific evidence
	Type of pyramidalis muscle dissection	Lack of procedure-specific evidence
	Rectus muscle re-approximation	Limited procedure-specific evidence

TAP, transversus abdominis plane block.

Surgical techniques that have been shown to be beneficial and are therefore recommended include Joel-Cohen incision and avoidance of peritoneum closure. Using abdominal binders postoperatively is recommended with sufficient procedure-specific evidence being identified. Analgesic adjuncts such as listening to music via headphones and use of transcutaneous electrical nerve stimulation may be associated with improved pain relief and are recommended when available.

Although pre-operative gabapentinoids were recommended previously, they are no longer recommended despite positive studies of their benefits due

to concerns about side-effects such as sedation and respiratory depression [156]. Furthermore, it is not clear if gabapentinoids add to our current recommendations of basic analgesia, i.v. dexamethasone and regional analgesia.

Several intra-operative interventions are not recommended due to inconsistent or limited or lack of procedure-specific evidence and/or concerns of side-effects (Table 2). For example, intra-operative dexmedetomidine infusion has been shown to provide improved postoperative pain relief; however, it is not recommended because its benefits on top of basic analgesia remain unknown, and due to concerns of side-

effects including hypotension and bradycardia which can be prolonged and might impede ambulation [157]. Similarly, although, a sub-anaesthetic dose of i.v. ketamine has demonstrated positive effects on postoperative pain scores [158], it is not recommended because its benefits over basic analgesia are unknown, and concerns of side-effects such as hallucinations that might impair the recollection of the birth experience and mother-child bonding [158,159].

Intrathecal or epidural administration of buprenorphine, hydromorphone, midazolam,  $\alpha$ 2-adrenergic agonists, neostigmine and ketamine has been reported to prolong the analgesic duration of morphine. However, they cannot be recommended due to inconsistent procedure-specific evidence and due to the potential side-effects such as hypotension or sedation. Additionally, in most studies, hypotension occurs as frequently as sedation. Peritoneal instillation of local anaesthetics cannot be recommended due to a lack of procedure-specific evidence. Similarly, topical local anaesthetic cream application is not recommended due to a lack of procedure-specific evidence.

The limitations of this review are related to those of the included studies. There was considerable heterogeneity between studies with regard to dosing regimens and route of administration as well as the timing of pain assessments. The small size of most studies makes it impossible to draw conclusions about the safety profile of an individual intervention. In the majority of included studies, the analgesic intervention was not evaluated against an optimised multimodal analgesic regimen. Moreover, measuring just pain scores and/or opioid consumption is not sufficient and more comprehensive, patient-centred tools to assess pain relief and functionality would better reflect day-to-day clinical practice but are unfortunately poorly reported in the literature. Also, because most studies include healthy, full-term parturients, our recommendations may not be applicable to parturients with co-existing medical conditions such as morbid obesity, chronic pain as well as preterm delivery. Furthermore, the PROSPECT methodology uses a minimal clinically important difference in pain scores of 1/10. However, this difference has never been validated in obstetric patients.

Future adequately powered studies should assess the effects of analgesic interventions not only on pain, opioid consumption, opioid-related adverse events and complications associated with the intervention, but also outcome measures such as time to ambulation, hospital stay and the occurrence of chronic pain or chronic opioid consumption. Furthermore, it is necessary to examine the

influence of analgesic interventions on patient-reported outcomes such as mother-child bonding, breast feeding ability, time to ambulation and return to activities of daily living. Validated scoring tools such as the quality of recovery-11 are useful metrics that should be considered.

In summary, this review has identified an analgesic regimen that can be used for optimal pain management after elective caesarean section performed under neuraxial anaesthesia. A combination of basic analgesics such as paracetamol; NSAIDs or cyclo-oxygenase-2-selective inhibitors; and i.v. dexamethasone, along with a local/regional analgesic techniques (e.g. intrathecal morphine 50–100  $\mu$ g or diamorphine 300  $\mu$ g); local anaesthetic infiltration with or without a field blocks such as ilio-inguinal and iliohypogastric nerves blocks or fascial plane blocks (e.g. TAP, quadratus lumborum or ESP blocks) are recommended. However, the benefits of local and regional analgesic techniques are not apparent with the use of intrathecal morphine or diamorphine. Analgesic adjuncts such as listening to music via headphones and transcutaneous electrical nerve stimulation may be used when available. Several aspects of the surgical technique clearly yield positive analgesic effects after caesarean section including the Joel-Cohen incision, non-closure of the peritoneum and the use of abdominal binders. The PROSPECT recommendation for postoperative analgesia after caesarean section has established a multimodal pre-, intra- and postoperative analgesic strategy which combined with certain surgical approaches and adjuvant techniques may provide excellent analgesia.

## Acknowledgements

This PROSPECT recommendation is supported by the Obstetric Anaesthetists' Association. PROSPECT is supported by an unrestricted grant from the European Society of Regional Anaesthesia and Pain Therapy. In the past, PROSPECT has received unrestricted grants from Pfizer Inc. New York, NY, USA and Grunenthal, Aachen, Germany. GJ has received honoraria from Baxter and Pacira Pharmaceuticals. MVdV has received honoraria from Sintetica, Grunenthal, Vifor Pharma, MSD, Nordic Pharma, Janssen Pharmaceuticals, Heron Therapeutics and Aquettant. EP-Z has received honoraria from Mundipharma, Grunenthal, MSD, Janssen-Cilag GmbH, Fresenius Kabi and AcelRx. No other external funding or competing interests declared.

## References

1. Gamez BH, Habib AS. Predicting severity of acute pain after Caesarean delivery: a narrative review. *Anesthesia and Analgesia* 2018; **126**: 1606–14.

2. Kainu JP, Sarvela J, Tiippana E, Halmesmaki E, Korttila KT. Persistent pain after caesarean section and vaginal birth: a cohort study. *International Journal of Obstetric Anesthesia* 2010; **19**: 4–9.
3. PROSPECT. Recommendations for postoperative analgesia after caesarean section 2014. [www.postoppain.org](http://www.postoppain.org) (accessed 04/11/2020).
4. Huang J, Cao C, Nelson G, Wilson RD. A review of enhanced recovery after surgery principles used for scheduled Caesarean delivery. *Journal of Obstetrics and Gynecology* 2019; **41**: 1775–88.
5. Joshi GP, Schug SA, Kehlet H. Procedure-specific pain management and outcome strategies. *Best Practice and Research Clinical Anaesthesiology*. 2014; **28**: 191–201.
6. Lee B, Schug SA, Joshi GP, Kehlet H. Procedure-specific pain management (PROSPECT) – an update. *Best Practice and Research Clinical Anaesthesiology* 2018; **32**: 101–11.
7. Joshi GP, Kehlet H. Guidelines for perioperative pain management: need for re-evaluation. *British Journal of Anaesthesia* 2017; **119**: 703–6.
8. Joshi GP, Van de Velde M, Kehlet H. Development of evidence-based recommendations for procedure-specific pain management: procedure-specific pain management (PROSPECT) methodology. *Anaesthesia* 2019; **74**: 1298–304.
9. Towers CV, Shelton S, van Nes J, et al. Preoperative caesarean delivery intravenous acetaminophen treatment for postoperative pain control: a randomized double-blind placebo control trial. *American Journal of Obstetrics and Gynecology* 2018; **218**: 353. e1–4.
10. Mahajan L, Mittal V, Gupta R, Chhabra H, Vidhan J, Kaur A. Study to compare the effect of oral, rectal and intravenous infusion of paracetamol for postoperative analgesia in women undergoing Cesarean section under spinal anesthesia. *Anesthesia Essays and Researches* 2017; **11**: 594–8.
11. Altenau B, Crisp CC, Devaiah CG, Lambers DS. Randomized controlled trial of intravenous acetaminophen for postcesarean delivery pain control. *American Journal of Obstetrics and Gynecology* 2017; **217**: 362. e1–6.
12. Bernstein J, Spitzer Y, Ohaegbulam K, Reddy S, Song J, Romanelli E, Nair S. The analgesic efficacy of IV acetaminophen for acute postoperative pain in C-section patients: a randomized, double-blind, placebo-controlled study. *Journal of Maternal-Fetal and Neonatal Medicine* 2020. Epub 10 March. <https://doi.org/10.1080/14767058.2020.1735337>.
13. Zeng AM, Nami NF, Wu CL, Murphy JD. The analgesic efficacy of nonsteroidal anti-inflammatory agents (NSAIDs) in patients undergoing cesarean deliveries, a meta-analysis. *Regional Anesthesia and Pain Medicine* 2016; **41**: 763–72.
14. Mkontwana N, Novikova N. Oral analgesia for relieving post-caesarean pain. *Cochrane Database of Systematic Reviews* 2015; **3**: CD010450.
15. Inthigood N, Lertbunnaphong JA. Efficacy of a single 40-mg intravenous dose of parecoxib for postoperative pain control after elective cesarean delivery: a double-blind randomized placebo controlled trial. *Journal of Obstetrics and Gynecology Research* 2017; **43**: 92–9.
16. Mahdavi A, Telkabadi Z, Aleyasin A, Hosseini MA, Safdarian L, Momenzadeh A. Comparison of morphine suppository and diclofenac suppository for pain management after elective caesarean section. *Acta Medicine Iran* 2016; **54**: 709–12.
17. Thippeswamy T, Krishnaswamy B, Begalorkar GM, Mariyappa N. Comparison of efficacy and safety of intramuscular piroxicam and tramadol for postoperative pain in patients undergoing Caesarean delivery. *Journal of Clinical and Diagnostic Research* 2016; **10**: FC01.
18. Khezri MB, Mosallaei MA, Ebtehaj M, Mohammadi N. Comparison of preemptive effect of ketorolac versus meperidine on postoperative shivering and pain in patients undergoing caesarean section under spinal anesthesia: a prospective, randomized, double-blind study. *Caspian Journal Internal Medicine* 2018; **9**: 151–7.
19. Olajetu SO, Adenekan AT, Olufolabi AJ, et al. Pentazocine versus pentazocine with rectal diclofenac for postoperative pain relief after caesarean section – a double blind randomized placebo controlled trial in a low resource area. *Middle East Journal of Anesthesia* 2016; **23**: 443–8.
20. Monks DT, Hoppe DW, Downey K, Shah V, Bernstein P, Carvalho JCA. A perioperative course of gabapentin does not produce a clinically meaningful improvement in analgesia after caesarean delivery. *Anesthesiology* 2015; **123**: 320–6.
21. El Kenany S, El Tahan MR. Effect of preoperative pregabalin on post-caesarean delivery analgesia: a dose-response study. *International Journal of Obstetric Anesthesia* 2016; **26**: 24–31.
22. Najafi Anaraki A, Mirzaei K. The effect of gabapentin versus intrathecal fentanyl on postoperative pain and morphine consumption in caesarean delivery: a prospective, randomized, double-blind study. *Archives Gynecology and Obstetrics* 2014; **290**: 47–52.
23. Khezri MB, Nasseh N, Soltanian G. The comparative preemptive analgesic efficacy of addition of vitamin B complex to gabapentin versus gabapentin alone in women undergoing cesarean section under spinal anesthesia: a prospective randomized double-blind study. *Medicine* 2017; **96**: 15.
24. Felder L, Saccone G, Scutto S, et al. Perioperative gabapentin and post cesarean pain control: a systematic review and meta-analysis of randomized controlled trials. *European Journal of Obstetrics and Gynecology and Reproductive Biology* 2019; **233**: 98–106.
25. Habibi A, Alipour A, Baradari AG, Gholinataj A, Habibi MR, Peivandi S. The effect of adding lidocaine to patient controlled analgesia with morphine on pain intensity after caesarean section with spinal anaesthesia: a double-blind, randomized, clinical trial. *Open Access Macedonian Journal of Medical Sciences* 2019; **7**: 1946–50.
26. Rahmanian M, Leysi M, Hemmati AA, Mirmohammadkhani M. The effect of low-dose intravenous ketamine on postoperative pain following Cesarean section with spinal anesthesia: a randomized clinical trial. *Oman Medical Journal* 2015; **30**: 11–6.
27. Wang J, Xu Z, Feng Z, Ma R, Zhang X. Impact of ketamine on pain management in Cesarean section: A systematic review and meta-analysis. *Pain Physician* 2020; **23**: 135–48.
28. Nie Y, Tu W, Shen X, et al. Dexmedetomidine added to sufentanil patient-controlled intravenous analgesia relieves the postoperative pain after cesarean delivery: a prospective randomized controlled multicenter study. *Nature Scientific Reports* 2018; **8**: 9952.
29. Shalu PS, Ghodki PS. To study the efficacy of intravenous dexamethasone in prolonging the duration of spinal anesthesia in elective cesarean section. *Anesthesia Essays and Researches* 2017; **11**: 321–5.
30. Ituk U, Thenuwara K. The effect of a single intraoperative dose of intravenous dexamethasone 8 mg on post-caesarean delivery analgesia: a randomized controlled trial. *International Journal of Obstetric Anesthesia* 2018; **35**: 57–63.
31. Maged AM, Deeb WS, Elbaradie S, et al. Comparison of local and intravenous dexamethasone on postoperative pain and recovery after caesarean section. a randomized controlled trial. *Taiwanese Journal of Obstetrics and Gynecology* 2018; **57**: 346–50.
32. Edomwonyi NP, Osazuwa MO, Iribhogbe OI, Esangbedo SE. Postoperative analgesia using bupivacaine wound infiltration with intravenous tramadol or dexamethasone following obstetric spinal anaesthesia. *Nigerian Journal of Clinical Practice* 2019; **20**: 1584–9.

33. Niklasson B, Arnelo C, Georgsson Ohman S, Segerdahl M, Blanck A. Oral oxycodone for pain after caesarean section: a randomized comparison with nurse administered IV morphine in a pragmatic study. *Scandinavian Journal of Pain* 2015; **7**: 17–24.
34. Chi X, Li M, Mei W, Liao M. Comparison of patient-controlled intravenous analgesia with sufentanil versus tramadol in post-caesarean section pain management and lactation after general anesthesia – a prospective, randomized, double-blind, controlled study. *Journal of Pain Research* 2017; **10**: 1521–7.
35. Nie JJ, Sun S, Huang SQ. Effect of oxycodone patient-controlled intravenous analgesia after cesarean section: a randomized controlled study. *Journal of Pain Research* 2017; **10**: 2649–55.
36. Zhu J, Xu C, Wang X, Shi W. Comparison of the analgesic effects of dezocine, tramadol and butorphanol after cesarean section. *Pakistanian Journal of Pharmacological Sciences* 2018; **31**: 2191–5.
37. Cai Q, Gong H, Fan M, Chen W, Cai L. The analgesic effect of tramadol combined with buorphanol on uterine cramping pain after repeat caesarean section: a randomized, controlled, double-blind study. *Journal of Anesthesia* 2020; **34**: 825–33.
38. Duan G, Bao X, Yang G, et al. Patient-controlled intravenous tramadol versus patient-controlled intravenous hydromorphone for analgesia after secondary cesarean delivery: a randomized controlled trial to compare analgesic, anti-anxiety and anti-depression effects. *Journal of Pain Research* 2019; **12**: 49–59.
39. Ffrench-O'Carroll R, Steinhäuser H, Duff S, et al. A randomized controlled trial comparing tapentadol with oxycodone in non-breastfeeding women post-elective cesarean section. *Current Medical Research and Opinion* 2019; **35**: 975–81.
40. Makela K, Palomaki O, Pokkinen S, Yli-Hankala A, Helminen M, Uotila J. Oral versus patient-controlled intravenous administration of oxycodone for pain relief after cesarean section. *Archives of Gynecology and Obstetrics* 2019; **300**: 903–9.
41. Sultan P, Halpern SH, Pushpanathan E, Patel S, Carvalho B. The effect of intrathecal morphine dose on outcomes after elective Cesarean delivery: a meta-analysis. *Anesthesia and Analgesia* 2016; **123**: 154–64.
42. Berger JS, Gonzalez A, Hopkins A, et al. Dose-response of intrathecal morphine when administered with intravenous ketorolac for post-caesarean analgesia: a two-center, prospective, randomized, blinded trial. *International Journal of Obstetric Anesthesia* 2016; **28**: 3–11.
43. Booth JL, Harris LC, Eisenach JC, Pan PH. A randomized controlled trial comparing two multimodal analgesic techniques in patients predicted to have severe pain after Cesarean delivery. *Anesthesia and Analgesia* 2016; **122**: 1114–9.
44. Kaufner L, Heimann S, Zander D, et al. Neuraxial anesthesia for pain control after cesarean section: a prospective randomized trial comparing three different neuraxial techniques in clinical practice. *Minerva Anestesiologica* 2015; **82**: 514–24.
45. Carvalho B, Sutton CD, Kowalczyk JJ, Flood PD. Impact of patient choice for different analgesic protocols on opioid consumption: a randomized prospective trial. *Regional Anesthesia and Pain Medicine* 2019; **44**: 578–85.
46. Carvalho B, Mirza F, Flood P. Patient choice compared with no choice of intrathecal morphine dose for caesarean analgesia: a randomized clinical trial. *British Journal of Anaesthesia* 2017; **118**: 762–71.
47. Sharpe EE, Molitor RJ, Arendt KW, et al. Intrathecal morphine versus intrathecal hydromorphone for analgesia after caesarean delivery. *Anesthesiology* 2020; **132**: 1382–91.
48. Allen TK, Mishriky BM, Klinger RY, Habib AS. The impact of neuraxial clonidine on postoperative analgesia and perioperative adverse effects in women having elective Cesarean section – a systematic review and meta-analysis. *British Journal of Anaesthesia* 2018; **120**: 228–40.
49. Fernandes HS, Bliancherene F, Vago TM, et al. Clonidine effect on pain after Cesarean delivery: a randomized controlled trial of different routes of administration. *Anesthesia and Analgesia* 2018; **127**: 165–70.
50. Carvalho FAE, Tenorio SB, Shiohara FT, Maria LR, Mota A. Randomized study of postcesarean analgesia with intrathecal morphine alone or combined with clonidine. *Journal of Clinical Anesthesia* 2016; **33**: 395–402.
51. Khezri MB, Rezaei M, Reihany MD, Javadi EHS. Comparison of postoperative analgesic effect of intrathecal clonidine and fentanyl added to bupivacaine in patients undergoing Cesarean section: a prospective randomized double-blind study. *Pain Research and Treatment* 2014; **2014**: 513628.
52. Yousef AA, Salem HA, Moustafa MZ. Effect of mini-dose epidural dexmedetomidine in elective cesarean section using combined spinal-epidural anesthesia: a randomized double-blind controlled study. *Journal of Anesthesia* 2015; **29**: 708–14.
53. Qi X, Chen D, Li G, et al. Comparison of intrathecal dexmedetomidine with morphine as adjuvants in Cesarean sections. *Biology Pharmacology Bulletin* 2016; **39**: 1455–60.
54. Bi YH, Cui XG, Zhang RQ, Song CY, Zhang YZ. Low dose of dexmedetomidine as an adjuvant to bupivacaine in cesarean surgery provides better intraoperative somato-visceral sensory block characteristics and postoperative analgesia. *Oncotarget* 2017; **8**: 63587–95.
55. Bi YH, Wu JM, Zhang YZ, Zhang RQ. Effect of different doses of intrathecal dexmedetomidine as an adjuvant combined with hyperbaric ropivacaine in patients undergoing caesarean section. *Frontiers in Pharmacology* 2020; **11**: 342.
56. Yang Y, Song C, Song C, Li C. Addition of dexmedetomidine to epidural morphine to improve anesthesia and analgesia for caesarean section. *Experimental and Therapeutic Medicine* 2020; **19**: 1747–54.
57. Mostafa MF, Herdan R, Fathy GM, et al. Intrathecal dexmedetomidine versus magnesium sulphate for postoperative analgesia and stress response after caesarean delivery: randomized controlled double-blind study. *European Journal of Pain* 2020; **24**: 182–91.
58. Weigl W, Bierylo A, Wielgus M, et al. Analgesic efficacy of intrathecal fentanyl during the period of highest analgesic demand after caesarean section. *Medicine* 2016; **95**: e3827.
59. Weigl W, Bierylo A, Wielgus M, et al. Perioperative analgesia after intrathecal fentanyl and morphine or morphine alone for caesarean section. *Medicine* 2017; **96**: e8892.
60. Ravindran R, Sajid B, Ramadas KT, Susheela I. Intrathecal hyperbaric bupivacaine with varying doses of buprenorphine for postoperative analgesia after Cesarean section: a comparative study. *Anesthesia Essays and Researches* 2017; **11**: 952–7.
61. Yang M, Wang L, Chen H, Tang Y, Chen X. Postoperative analgesic effects of different doses of epidural hydromorphone coadministered with ropivacaine after Cesarean section: a randomized controlled trial. *Pain Research and Management* 2019; **2019**: 99054538.
62. Wang SC, Pan PT, Chiu HY, Huang CJ. Neuraxial magnesium sulfate improves postoperative analgesia in Cesarean section delivery women: a meta-analysis of randomized controlled trials. *Asian Journal of Anesthesiology* 2017; **55**: 56–67.
63. Abdollahpour A, Azadi R, Bandari R, Mirmohammadkhani M. Effects of adding midazolam and sufentanil to intrathecal bupivacaine on analgesia quality and postoperative complications in elective Cesarean section. *Anesthesiology and Pain Medicine* 2015; **5**: e23565.

64. Dodawad R, Sumalatha GB, Pandarpurkar S, Jajee P. Intrathecal midazolam as an adjuvant in pregnancy-induced hypertensive patients undergoing an elective Caesarean section: a clinical comparative study. *Anesthesiology and Pain Medicine* 2016; **6**: e38550.
65. Khezri MB, Tahaei E, Atlasbaf AH. Comparison of postoperative analgesic effect of intrathecal ketamine and fentanyl added to bupivacaine in patients undergoing cesarean section: a prospective randomized double-blind study. *Middle East Journal of Anesthesia* 2016; **23**: 427–36.
66. Basuni AS. Addition of low-dose ketamine to midazolam and low-dose bupivacaine improves hemodynamics and postoperative analgesia during spinal anesthesia for cesarean section. *Journal of Anaesthesiology and Clinical Pharmacology* 2016; **32**: 44–8.
67. Cossu AP, De Giudici LM, Piras D, et al. A systematic review of the effects of adding neostigmine to local anesthetics for neuraxial administration in obstetric anesthesia and analgesia. *International Journal of Obstetric Anesthesia* 2015; **24**: 237–46.
68. Hussien RM, Rabie AH. Sequential intrathecal injection of fentanyl and hyperbaric bupivacaine at different rates: does it make a difference? A randomized controlled trial. *Korean Journal of Anesthesiology* 2019; **72**: 150–5.
69. Patel R, Carvalho JCA, Downey K, Kanczuk M, Bernstein P, Siddiqui N. Intraperitoneal installation of lidocaine improves postoperative analgesia at cesarean delivery: a randomized, double-blind, placebo-controlled trial. *Anesthesia and Analgesia* 2017; **124**: 554–9.
70. Grosse-Steffen T, Kramer M, Tuschy B, Weiss C, Sutterlin M, Berlit S. Topic anaesthesia with a eutectic mixture of lidocaine/prilocaine cream after elective caesarean section: a randomized, placebo-controlled trial. *Archives of Gynecology and Obstetrics* 2017; **296**: 771–6.
71. Larsen KR, Kristensen BB, Rasmussen MA, et al. Effect of high volume systematic local infiltration analgesia in Caesarean section: a randomized, placebo-controlled trial. *Acta Anaesthesiologica Scandinavica* 2015; **59**: 632–9.
72. Ghenaee MM, Rahmani S, Jafarabadi M. Local lidocaine 2% in postoperative pain management in Caesarean delivery. *Journal of Family and Reproductive Health* 2015; **9**: 19–21.
73. Nasir F, Sohail I, Sadiq H, Habib M. Local wound infiltration with ropivacaine for postoperative pain control in Caesarean section. *Cureus* 2019; **11**: e5572.
74. Reinikainen M, Syvaaja S, Hara K. Continuous wound infiltration with ropivacaine for analgesia after caesarean section: a randomized placebo-controlled trial. *Acta Anaesthesiologica Scandinavica* 2014; **58**: 973–9.
75. Jolly C, Jathières F, Keita H, Jaouen E, Guyot B, Torre A. Cesarean analgesia using levobupivacaine continuous wound infiltration: a randomized trial. *European Journal of Obstetrics and Gynecology and Reproductive Biology* 2015; **194**: 125–30.
76. Lalmand M, Wilwerth M, Fils JF, Van der Linden P. Continuous ropivacaine subfascial wound infusion compared with intrathecal morphine for postcesarean analgesia: a prospective, randomized controlled double-blind study. *Anesthesia and Analgesia* 2017; **125**: 907–12.
77. Adesope O, Ituk U, Habib AS. Local anaesthetic wound infiltration for postcaesarean section analgesia. *European Journal of Anaesthesiology* 2016; **33**: 731–42.
78. Thomas D, Panneerselvam S, Kundra P, Rudingwa P, Sivakumar RK, Dorairajan G. Continuous wound infiltration of bupivacaine at two different anatomical planes for caesarean analgesia – a randomized clinical trial. *Indian Journal of Anaesthesia* 2019; **63**: 437–43.
79. Wagner-Kovacec J, Povalej-Brzan P, Mekis D. Efficacy of continuous in-wound infusion of levobupivacaine and ketorolac for post-caesarean section analgesia: a prospective, randomized, double-blind, placebo-controlled trial. *BMC Anesthesiology* 2018; **18**: 165.
80. Barney EZ, Pedro CD, Gamez BH, Fuller ME, Dominguez JE, Habib AS. Ropivacaine and ketorolac wound infusion for post-cesarean delivery analgesia. *Obstetrics and Gynecology* 2020; **135**: 427–35.
81. Kundra S, Singh RM, Singh G, Singh T, Jarewal V, Katyal S. Efficacy of magnesium sulphate as an adjunct to ropivacaine in local infiltration for postoperative pain following lower segment caesarean section. *Journal of Clinical and Diagnostic Research* 2016; **10**: UC18–UC22.
82. Bhardwaj S, Devgan S, Sood D, Katyal S. Comparison of local wound infiltration with ropivacaine alone or ropivacaine plus dexmedetomidine for postoperative pain relief after lower segment caesarean section. *Anesthesia Essays and Researches* 2017; **11**: 940–5.
83. Liu MW, Li TKT, Lui F, Ong CYT. A randomized, controlled trial of rectus sheath bupivacaine and intrathecal morphine, vs intrathecal bupivacaine and morphine after Caesarean section. *Anaesthesia* 2017; **72**: 1225–9.
84. Triyasunant N, Chinachoti T, Duangburong S. Direct field block with 40 mL of 0.125% bupivacaine in conjunction with intrathecal morphine for analgesia after Cesarean section: a randomized controlled trial. *Journal of the Medical Association of Thailand* 2015; **98**: 1001–9.
85. McKeen DM, George RB, Boyd JC, Allen VM, Pink A. Transversus abdominis plane block does not improve early or late pain outcomes after Cesarean delivery: a randomized controlled trial. *Canadian Journal of Anesthesia* 2014; **61**: 631–40.
86. Fusco P, Cofini V, Petrucci E, et al. Transversus abdominis plane block in the management of acute postoperative pain syndrome after Caesarean section: a randomized controlled clinical trial. *Pain Physician* 2016; **19**: 583–91.
87. Jadon A, Jain P, Chakraborty S, et al. Role of ultrasound guided transversus abdominis plane block as a component of multimodal analgesic regimen for lower segment caesarean section: a randomized double blind clinical study. *BMC Anesthesiology* 2018; **18**: 53.
88. Kupiec A, Zwierzchowski J, Kowal-Janicka J, et al. The analgesic efficiency of transversus abdominis plane (TAP) block after caesarean section. *Ginekologia Polska* 2018; **89**: 420–3.
89. Kakade A, Wagh G. Evaluate the feasibility of surgical transversus abdominis plane block for postoperative analgesia after Cesarean section. *Journal of Obstetrics and Gynecology of India* 2019; **69**: 330–3.
90. Faiz SHR, Alebouyeh MR, Derakhshan P, Imani F, Rahimzadeh P, Ashtiani MG. Comparison of ultrasound-guided posterior transversus abdominis plane block and lateral transversus abdominis plane block for postoperative pain management in patients undergoing caesarean section: a randomized double-blind clinical trial. *Journal of Pain Research* 2017; **11**: 5–9.
91. Narasimhulu DM, Scharfman L, Minkoff H, George B, Homel P, Tyagaraj K. A randomized trial comparing surgeon-administered intraoperative transversus abdominis plane block with anesthesiologist-administered transcutaneous block. *International Journal of Obstetric Anesthesia* 2018; **35**: 26–32.
92. Gupta A, Gupta A, Yadav N. Effect of dexamethasone as an adjuvant to ropivacaine on duration and quality of analgesia in ultrasound-guided transversus abdominis plane block in patients undergoing lower segment caesarean section – a prospective, randomized, single-blinded study. *Indian Journal of Anaesthesia* 2019; **63**: 469–74.
93. John R, Ranjan RV, Ramachandran TR, George SK. Analgesic efficacy of transverse abdominal plane block after elective cesarean delivery – bupivacaine with fentanyl versus

- bupivacaine alone: a randomized, double-blind controlled clinical trial. *Anesthesia Essays and Researches* 2017; **11**: 181–4.
94. Singh R, Kumar N, Jain A, Joy S. Addition of clonidine to bupivacaine in transversus abdominis plane block prolongs postoperative analgesia after caesarean section. *Journal of Anaesthesiology Clinical Pharmacology* 2016; **32**: 501–4.
  95. Acharya R, Baksi R, Mohapatra P. Comparative analysis of duration of postoperative analgesia between levobupivacaine and levobupivacaine with clonidine after ultrasound-guided transversus abdominis plane block in patients undergoing lower segment caesarean section. *Anesthesia Essays and Researches* 2018; **12**: 943–8.
  96. Parameswari AR, Udayakumar P. Comparison of efficacy of bupivacaine with dexmedetomidine versus bupivacaine alone for transversus abdominis plane block for postoperative analgesia in patients undergoing elective caesarean section. *Journal of Obstetrics and Gynecology of India* 2018; **62**: 98–103.
  97. Canakci E, Gultekin A, Cebeci Z, Hanedan B, Kilinc A. The analgesic efficacy of transverse abdominis plane block versus epidural block after caesarean delivery: Which one is effective? TAP block? Epidural block? *Pain Research and Management* 2018; **2018**: 3562701.
  98. Jarraya A, Zghal J, Abidi S, Smaoui M, Kolsi K. Subarachnoid morphine versus TAP blocks for enhanced recovery after caesarean section delivery: a randomized controlled trial. *Anaesthesia, Critical Care and Pain Medicine* 2016; **35**: 401–3.
  99. Dereu D, Savoldelli GL, Mercier Y, Combescure C, Mathivon S, Rehberg B. The impact of a transversus abdominis plane block including clonidine vs intrathecal morphine on nausea and vomiting after caesarean section. *European Journal of Anaesthesiology* 2019; **36**: 575–82.
  100. Kwikiriza A, Kiwanuka JK, Firth PG, Hoeft MA, Modest VE, Ttendo SS. The analgesic effects of intrathecal morphine in comparison with ultrasound-guided transversus abdominis plane block after caesarean section: a randomized controlled trial at a Ugandan regional referral hospital. *Anaesthesia* 2019; **74**: 167–73.
  101. Telnes A, Skogvoll E, Lonnée H. Transversus abdominis plane block vs wound infiltration in Caesarean section: a randomized controlled trial. *Acta Anaesthesiologica Scandinavica* 2015; **59**: 496–504.
  102. Klasen F, Bourgoin A, Antonini F, et al. Postoperative analgesia after caesarean section with transversus abdominis plane block or continuous infiltration wound catheter: A randomized clinical trial. TAP vs infiltration after caesarean section. *Anaesthesiology, Critical Care and Pain Medicine* 2016; **35**: 411–6.
  103. Tawfik MM, Mohamed YM, Elbadrawi RE, Abdelkhalik M, Mogadeh MM, Ezz HM. Transversus abdominis plane block versus wound infiltration for analgesia after caesarean delivery: a randomized controlled trial. *Anesthesia and Analgesia* 2017; **124**: 1291–7.
  104. Fusco P, Scimia P, Paladini G, et al. Transversus abdominis plane block for analgesia after Caesarean delivery. A systematic review. *Minerva Anestesiologica* 2015; **81**: 195–204.
  105. Champaneria R, Shah L, Wilson MJ, Daniels JP. Clinical effectiveness of transversus abdominis plane (TAP) blocks for pain relief after caesarean section: a meta-analysis. *International Journal of Obstetric Anesthesia* 2016; **28**: 45–60.
  106. Ng SC, Habib AS, Sodha S, Carvalho B, Sultan P. High-dose versus low-dose local anaesthetic for transversus abdominis plane block post-Caesarean delivery analgesia: a meta-analysis. *British Journal of Anaesthesia* 2018; **120**: 252–63.
  107. Staker JJ, Liu D, Church R, et al. A triple-blind, placebo-controlled randomized trial of the ilioinguinal-transversus abdominis plane (I-TAP) nerve block for elective caesarean section. *Anaesthesia* 2018; **73**: 594–602.
  108. Qian H, Zhang Q, Zhu P, et al. Ultrasound-guided transversus abdominis plane block using ropivacaine and dexmedetomidine in patients undergoing caesarean sections to relieve post-operative analgesia: a randomized controlled clinical trial. *Experimental and Therapeutic Medicine* 2020; **20**: 1163–8.
  109. Blanco R, Ansari T, Girgis E. Quadratus lumborum block for postoperative pain after caesarean section. *European Journal of Anaesthesiology* 2015; **32**: 812–8.
  110. Krohg A, Ullensvang K, Rosseland LA, Langesaeter E, Sauter AR. The analgesic effect of ultrasound-guided quadratus lumborum block after Caesarean delivery: a randomized clinical trial. *Anesthesia and Analgesia* 2018; **126**: 559–65.
  111. Hansen CK, Dam M, Steingrimsdottir GE, et al. Ultrasound-guided transmuscular quadratus lumborum block for elective caesarean section significantly reduces postoperative opioid request: a double-blind randomized trial. *Regional Anesthesia and Pain Medicine* 2019; **44**: 896–900.
  112. Mieszkowski MM, Mayzner-Zawadzka E, Tuyakov B, et al. Evaluation of the effectiveness of the quadratus lumborum block type I using ropivacaine in postoperative analgesia after a caesarean section – a controlled clinical study. *Ginekologie Polska* 2018; **89**: 89–96.
  113. Blanco R, Ansari T, Riad W, Shetty N. Quadratus lumborum block versus transversus abdominis plane block for postoperative pain after caesarean delivery. *Regional Anesthesia and Pain Medicine* 2016; **41**: 757–62.
  114. Verma K, Malawat A, Jethava D, Jethava DD. Comparison of transversus abdominis plane block and quadratus lumborum block for post-caesarean section analgesia: a randomized clinical trial. *Indian Journal of Anaesthesia* 2019; **63**: 820–6.
  115. Kang W, Lu D, Yang X, et al. Postoperative analgesic effects of various quadratus lumborum block approaches following caesarean section: a randomized controlled trial. *Journal of Pain Research* 2019; **12**: 2305–12.
  116. Tamura T, Yokota S, Ando M, Kubo Y, Nishiwaki K. A triple-blinded randomized trial comparing spinal morphine with posterior quadratus lumborum block after caesarean section. *International Journal of Obstetric Anesthesia* 2019; **40**: 32–8.
  117. Salama ER. Ultrasound-guided bilateral quadratus lumborum block vs. intrathecal morphine for postoperative analgesia after caesarean section: a randomized controlled trial. *Korean Journal of Anesthesiology* 2020; **73**: 121–8.
  118. Yoshida K, Tanaka S, Watanabe K, Obara S, Murakawa M. The effectiveness of the intramuscular quadratus lumborum block in postoperative analgesia after caesarean section with vertical incision: a randomized, double-blind placebo-controlled study. *Journal of Anesthesia* 2020; **34**: 849–56.
  119. Irwin R, Stanescu S, Buzaianu C, Rademan M, Roddy J, Gormley C, Tan T. Quadratus lumborum block for analgesia after caesarean section: a randomised controlled trial. *Anaesthesia* 2020; **75**: 89–95.
  120. Sultan P, Patel SD, Jadin S, Carvalho B, Halpern SH. Transversus abdominis plane block compared with wound infiltration for postoperative analgesia following Caesarean delivery: a systematic review and network meta-analysis. *Canadian Journal of Anesthesia* 2020; **67**: 1710–27.
  121. El-Boghdady K, Desai N, Halpern S, et al. Quadratus lumborum block versus transversus abdominis plane block for caesarean delivery: a systematic review and network meta-analysis. *Anaesthesia* 2020. Epub 4 July. <https://doi.org/10.1111/anae.15160>.
  122. Boules ML, Goda AS, Abdelhady MA, Abu El-Nour Abd El-Azeem SA, Hamed MA. Comparison of analgesic effect between erector spinae plane block and transversus abdominis plane block after elective caesarean section: a prospective randomized single-blind controlled study. *Journal of Pain Research* 2020; **13**: 1073–80.

123. Hamed MA, Yassin HM, Botros JM, Abdelhady MA. Analgesic efficacy of erector spinae plane block compared with intrathecal morphine after elective caesarean section: a prospective randomized controlled study. *Journal of Pain Research* 2020; **13**: 597–604.
124. Sato I, Iwasaki H, Luthe SK, Lida T, Kanda H. Comparison of intrathecal morphine with continuous patient-controlled epidural anesthesia versus intrathecal morphine alone for post-caesarean section analgesia: a randomized controlled trial. *BMC Anesthesiology* 2020; **20**: 138.
125. Chen SY, Liu FL, Cherng YG, et al. Patient-controlled epidural levobupivacaine with or without fentanyl for post-caesarean section pain relief. *Biomedical Research International* 2014; 965152.
126. Khooshideh M, Rostami SSL, Sheikh M, Yekta BG, Shahriari A. Pulsed electromagnetic fields for postsurgical pain management in women undergoing caesarean section. A randomized, double-blind, placebo-controlled trial. *Clinical Journal of Pain* 2017; **33**: 142–7.
127. Chakravarthy M, Prashanth A, George A. Evaluation of percutaneous electrical nerve stimulation of the auricle for relief of postoperative pain following caesarean section. *Medical Acupuncture* 2019; **31**: 281–8.
128. Bonnal A, Dehon A, Nagot N, Macioce V, Nogue E, Moreau E. Patient-controlled oral analgesia versus nurse-controlled parenteral analgesia after caesarean section: a randomized controlled trial. *Anaesthesia* 2016; **71**: 535–43.
129. Yefet E, Taha H, Salim R, et al. Fixed time interval compared with on-demand oral analgesia protocols for post-caesarean pain: a randomized controlled trial. *British Journal of Obstetrics and Gynaecology* 2017; **124**: 1063–70.
130. Farzaneh M, Abbasjahreni A, Saadatmand V, et al. Comparative effect of nature-based sounds intervention and headphones intervention on pain severity after Caesarean section: a prospective double-blind randomized trial. *Anesthesiology and Pain Medicine* 2019; **9**: e67835.
131. Kollmann M, Adrian L, Scheuchenegger A, et al. Early skin-to-skin contact after caesarean section: a randomized clinical pilot study. *PLoS One* 2017; **12**: e0168783.
132. Ghana S, Hakimi S, Mirghafourvand M, Abbasalizadeh F, Behnampour N. Randomized controlled trial of abdominal binders for postoperative pain, distress and blood loss after caesarean delivery. *International Journal of Gynecology and Obstetrics* 2017; **137**: 271–6.
133. Gustafson JL, Dong F, Duong J, Kuhlmann ZC. Elastic abdominal binders reduce caesarean pain postoperatively: a randomized controlled pilot trial. *Kansas Journal of Medicine* 2018; **11**: 48–53.
134. Karaca I, Ozturk M, Alay I, et al. Influence of abdominal binder usage after caesarean delivery on postoperative mobilization, pain and distress: a randomized controlled trial. *Eurasian Journal of Medicine* 2019; **51**: 214–8.
135. Sakinci M, Kuru O, Olgan S, et al. Dilatation of the cervix at non-labour caesarean section: does it improve the patients perception of pain postoperatively? *Journal of Obstetrics and Gynecology* 2015; **35**: 681–4.
136. Alalfy M, Yehia A, Samy A. Routine cervical dilatation at caesarean section and its influence on postoperative pain and complications in obese women: a double blind randomized controlled trial. *Journal of Maternal-Fetal and Neonatal Medicine* 2019; **2019**: 1651274.
137. Goymen A, Simsek Y, Ozdurak HI, et al. Effect of vaginal cleansing on postoperative factors in elective caesarean sections: a prospective, randomized controlled trial. *Journal of Maternal-Fetal and Neonatal Medicine* 2017; **30**: 442–5.
138. Gizzo S, Andrisani A, Noventa M, et al. Caesarean section: could different transverse abdominal incision techniques influence postpartum pain and subsequent quality of life? A systematic review. *PLoS One* 2015; **10**: e0114190.
139. Elbohotey AEH, Gomaa MF, Abdelaleim M, Abd-El-Gawad M, Elmarakby M. Diathermy versus scalpel in transverse abdominal incision in women undergoing repeated caesarean section: a randomized controlled trial. *Journal of Obstetrics and Gynaecology Research* 2015; **41**: 1541–6.
140. Yilmaz FY, Mathyk BA, Yildiz S, Yenigul NN, Saglam C. Postoperative pain and neuropathy after caesarean operation featuring blunt or sharp opening of the fascia: a randomized parallel group double-blind study. *Journal of Obstetrics and Gynaecology* 2018; **38**: 933–9.
141. Yapca OE, Topdagi YE, Ragip AA. Fetus delivery time in extraperitoneal versus transperitoneal caesarean section: a randomized trial. *Journal of Maternal-Fetal and Neonatal Medicine* 2020; **33**: 657–63.
142. Akhlagi F, Khazaie A, Jafaripour F. Comparing formation or non-formation of bladder flap at caesarean section on perioperative and postoperative complications: double-blind clinical trial. *Journal of Family and Reproductive Health* 2017; **11**: 152–8.
143. El-Khayat W, Elsharkawi M, Hassan A. A randomized controlled trial of uterine exteriorization versus in situ repair of the uterine incision caesarean delivery. *International Journal of Gynecology and Obstetrics* 2014; **127**: 163–6.
144. Zaphiratos V, George RB, Boyd JC, Habib AS. Uterine exteriorization compared with in situ repair for Caesarean delivery: a systematic review and meta-analysis. *Canadian Journal of Anesthesia* 2015; **62**: 1209–20.
145. Skret-Magierlo J, Soja P, Drozdowska A, et al. Two techniques of pyramidalis muscle dissection in Pfannenstiel incision for caesarean section. *Ginekologia Polska* 2015; **86**: 509–13.
146. Eken MK, Ozkaya E, Tarhan T, et al. Effects of closure versus non-closure of the visceral and parietalperitoneum at caesarean section: does it have any effect on postoperative vital signs? A prospective randomized study. *Journal of Maternal-Fetal and Neonatal Medicine* 2017; **30**: 922–6.
147. Omran EF, Meshaal H, Hassan SM, Dieb AS, Nabil H, Saad H. The effect of rectus muscle re-approximation at caesarean delivery on pain perceived after operation: a randomized control trial. *Journal of Maternal-Fetal and Neonatal Medicine* 2018; **32**: 3238–43.
148. Bamigboye AA, Hofmeyr GJ. Closure versus non-closure of the peritoneum at caesarean section: short- and long-term outcomes. *Cochrane Database of Systematic Reviews* 2014; **8**: CD000163.
149. Poursalehan S, Nesioonpour S, Akhondzadeh R, Mokmeli S. The effect of low-level laser on postoperative pain after elective caesarean section. *Anesthesia Pain Medicine* 2018; **8**: e84195.
150. Pires de Holanda AAM, Rocha de Sena KR, da Silva M, Filho E, Pegado R, Albuquerque Barbosa Cabral Micussi MT. Low-level laser therapy improves pain in postcaesarean section: a randomized clinical trial. *Lasers in Medical Science* 2020; **35**: 1095–102.
151. Maged AM, Mohesen MN, Elhalwagy A, et al. Subcuticular interrupted versus continuous skin suturing in elective caesarean section in obese women: a randomized controlled trial. *Journal of Maternal-Fetal and Neonatal Medicine* 2019; **32**: 4114–9.
152. Mackeen AD, Schuster M, Berghella V. Suture versus staples for skin closure after caesarean: a metaanalysis. *American Journal of Obstetrics and Gynecology* 2015; **212**(621): e1–10.
153. Wang H, Hong S, Teng H, Qiao L, Yin H. Subcuticular sutures versus staples for skin closure after caesarean delivery: a meta-analysis. *Journal of Maternal-Fetal and Neonatal Medicine* 2016; **29**: 3705–11.

154. Sharawi N, Carvalho B, Habib AS, Blake L, Jm M, Sultan P. A systematic review evaluating neuraxial morphine and diamorphine associated respiratory depression after Cesarean delivery. *Anesthesia and Analgesia* 2018; **127**: 1385–95.
155. NICE. Caesarean section. [CG132]. last updated September 2019. [www.nice.org.uk/guidance/cg132/](http://www.nice.org.uk/guidance/cg132/) (accessed 30/06/2020).
156. Cavalcante AN, Sprung J, Schroeder DR, Weingarten TN. Multimodal analgesic therapy with gabapentin and its association with postoperative respiratory depression. *Anesthesia and Analgesia* 2017; **125**: 141–6.
157. Grape S, Kirkham KR, Frauenknecht J, Albrecht E. Intra-operative analgesia with remifentanyl vs. dexmedetomidine: a systematic review and meta-analysis with trial sequential analysis. *Anaesthesia* 2019; **74**: 793–800.
158. Bauchat JR, Higgins N, Wojciechowski KG, McCarthy RJ, Toledo P, Wong CA. Low-dose ketamine with multimodal postcesarean delivery analgesia: a randomized controlled trial. *International Journal of Obstetric Anesthesia* 2011; **20**: 3–9.
159. Carvalho B, Butwick AJ. Postcesarean delivery analgesia. *Best Practice and Research Clinical Anaesthesiology* 2017; **31**: 69–79.

## Appendix 1

### PROSPECT Working Group

G. P. Joshi, E. Pogatzki-Zahn, M. Van de Velde, S. Schug, H. Kehlet, F. Bonnet, N. Rawal, A. Delbos, P. Lavand'homme, H.

Beloeil, J. Raeder, A. Sauter, E. Albrecht, P. Lirk, D. Lobo, S. Freys.

### Supporting Information

Additional supporting information may be found online via the journal website.

**Table S1.** Quality assessment and level of evidence assigned to the randomised trials included in the review for analgesia after caesarean section.

**Table S2.** Summary of key results from studies evaluating systemic analgesics, analgesics adjuncts, regional anaesthesia and surgical procedures used to support the recommended interventions in patients after caesarean section.

**Table S3.** Summary of key results from studies evaluating systemic analgesics, analgesics adjuncts, regional anaesthesia and surgical procedures used to support the interventions that are not recommended in patients undergoing caesarean section.